OncoLink
Abramson Cancer Center of the University of Pennsylvania

Vinblastine (Velban®, Alkaban AQ)

Pronounced: vin-BLAS-teen
Classification: Antimicrotubule Agent

About Vinblastine

Vinblastine is a member of the vinca alkaloid family of chemotherapy drugs. The drug works by interfering with cell division, which leaves the tumor unable to grow and spread. Vinblastine was developed from the Madagascar periwinkle plant.

How to Take Vinblastine

This medication is given through an IV (into a vein) by a trained professional. Specific dosage and dosing schedule depends on the person's size and the type of cancer being treated.

This medication is a "vesicant". This means that even when carefully and correctly administered by trained personnel, this drug may cause feeling of burning and pain. There is a risk that this drug may leak out of the vein at the injection site, resulting in tissue damage that can be severe. If the area of injection becomes red, swollen, or painful at anytime during or after the injection, notify your doctor or nurse immediately. Do not apply anything to the site unless instructed by your doctor or nurse.

Possible Side Effects of Vinblastine

There are a number of things you can do to manage the side effects of Vinblastine. Talk to your doctor or nurse about these recommendations. They can help you decide what will work best for you. These are some of the most common side effects:

Constipation

Vinblastine can cause serious constipation, abdominal pain and can even lead to a blockage or stoppage of the bowel (called paralytic ileus) if not treated promptly. There are several things you can do to prevent or relieve constipation. Include fiber in your diet (fruits and vegetables), drink 8-10 glasses of non-alcoholic fluids a day, and keep active. A stool softener once or twice a day may prevent constipation. If you do not have a bowel movement for 2-3 days, you should contact your healthcare team for suggestions to relieve the constipation.

Low White Blood Cell Count (Leukopenia or Neutropenia

White blood cells (WBC) are important for fighting infection. While receiving treatment, your WBC count can drop, putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4°), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

Tips to preventing infection:

- Washing hands, both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent mouth care.
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your doctor or nurse before scheduling dental appointments or procedures.
- Ask your doctor or nurse before you, or someone you live with, has any vaccinations.

For more suggestions, read the Neutropenia Tip Sheet.

Low Red Blood Cell Count (Anemia

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the red cell count is low, you may feel tired or weak. You should let your doctor or nurse know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion. Read the anemia tip sheet for more information.

Low Platelet Count (Thrombocytopenia

Platelets help your blood clot, so when the count is low you are at a higher risk of bleeding. Let your doctor or nurse know if you have any excess bruising or bleeding, including nose bleeds, bleeding gums or blood in your urine or stool. If the count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin®, Aleve®, Advil®, etc. as these can all increase the risk of bleeding. Unless your doctor tells you otherwise, you may take acetaminophen (Tylenol).

- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

Read the thrombocytopenia tip sheet for more information.

Nausea and/or Vomiting

Take anti-nausea medications as prescribed. If you continue to have nausea or vomiting, notify your doctor or nurse so they can help you manage this side effect. In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try antacids, (e.g. milk of magnesia, calcium tablets such as Tums), saltines, or ginger ale to lessen symptoms. Read the Nausea & Vomiting Tip Sheet for more suggestions.

Call your doctor or nurse if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

Fatigue

While on cancer treatment you may need to adjust your schedule to manage fatigue. Plan times to rest during the day and conserve energy for more important activities. Exercise can help combat fatigue; a simple daily walk with a friend can help. Talk to your healthcare team and see OncoLink's section on fatigue for helpful tips on dealing with this side effect.

Sexual and Reproductive Changes

This drug can affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and vaginal dryness. The desire for sex may decrease during treatment.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment, even if your menstrual cycle stops or you believe your sperm is affected. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team. See OncoLink's section on sexuality for helpful tips for dealing with these side effects.

Decrease in Appetite

Visit OncoLink's section on Nutrition for tips on dealing with this side effect. Ask your nurse about nutritional counseling services.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.

- You may experience a metallic taste or dislike foods or beverages that you liked before receiving chemotherapy. These symptoms can last up to several months.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products and fish without a strong smell.
- Flavor meat or fish by marinating it in sweet juices, sweet and sour sauce or dressings. Use seasonings like basil, oregano or rosemary. Bacon, ham and onion can also add flavor to vegetables.

Mouth Ulcers (Sores)

Certain cancer treatments can cause sores or soreness in your mouth and/or throat. Notify your doctor or nurse if your mouth, tongue, inside of your cheek or throat becomes white, ulcerated or painful. Performing regular mouth care can help prevent or manage mouth sores. If mouth sores become painful, your doctor or nurse can recommend a pain reliever.

- Brush with a soft-bristle toothbrush or cotton swab twice a day.
- Avoid mouthwashes that contain alcohol. A baking soda and/or salt warm water mouth rinse (2 level teaspoons of baking soda or 1 level teaspoon salt in an eight ounce glass of warm water) is recommended 4 times daily.
- If your mouth becomes dry, eat moist foods, drink plenty of fluids (6-8 glasses), and suck on sugarless hard candy.
- Avoid smoking and chewing tobacco, drinking alcoholic beverages and citrus juices.

Read the mouth ulcer tip sheet for more information.

Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become thin, brittle, or may fall out. This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun. Read more on alopecia.

Neurotoxicity

This is a toxicity that affects the nerves. The most common affect is called peripheral neuropathy, which causes a numbness or tingling of the hands and feet, which can get progressively worse with subsequent doses. You should let your healthcare provider know if you experience numbness or tingling in the hands and feet, as they may need to change the doses of your medication. See OncoLink's section on peripheral neuropathy for tips on dealing with this side effect.

The vinca alkaloid chemotherapies are known to cause neuropathy, but vinblastine can also cause neurologic toxicity that presents as mental depression, headache, malaise, dizziness and seizures. It can also cause toxicity of the cranial nerves, which affects

the vocal cords (changes in voice), eyes (visual changes) or facial nerves (drooping of the face or mouth). If you notice any of these problems, notify your healthcare team right away.

Other Side Effects

Patients can develop severe pain in the jaw soon after the first treatment with vinblastine. Your healthcare team can recommend a pain reliever for this problem, which will go away in time. Urinary retention can develop. If you notice you are unable to urinate, you should call your healthcare team right away or go to the emergency room to be evaluated.

Ifosfamide (Ifex®)

Pronounced: eye-FOSS-fuh-MIDE
Classification: Alkylating Agent

About Ifosfamide

Ifosfamide exerts its anti-cancer affect by a process called alkylation. Alkylation damages the DNA of cells, which prevents them from dividing and causes them to die. Since cancer cells, in general, divide faster and with less error correcting than healthy cells, cancer cells are more sensitive to this damage.

How to Take Ifosfamide

Ifosfamide is given by intravenous (into a vein) infusion. The dosage and schedule is determined by the person's size and type of cancer. It can be given alone or with other drugs.

Possible Side Effects of Ifosfamide

There are a number of things you can do to manage the side effects of Ifosfamide. Talk to your doctor or nurse about these recommendations. They can help you decide what will work best for you. These are some of the most common side effects:

Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your WBC count can drop, putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4 F), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

Tips to preventing infection:

- Washing hands, both yours and your visitors, is the best way to prevent the

spread of infection.

- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever or cough or live with someone with these symptoms).
- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent mouth care.
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your doctor or nurse before scheduling dental appointments or procedures.
- Ask your doctor or nurse before you, or someone you live with, has any vaccinations.

For more suggestions, read the Neutropenia Tip Sheet.

Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the red cell count is low, you may feel tired or weak. You should let your doctor or nurse know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion. Read the anemia tip sheet for more information.

Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the count is low you are at a higher risk of bleeding. Let your doctor or nurse know if you have any excess bruising or bleeding, including nosebleeds, bleeding gums or blood in your urine or stool. If your platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor to shave (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin®, Aleve®, Advil®, etc. as these can all increase the risk of bleeding. Unless your healthcare team tells you otherwise, you may take acetaminophen (Tylenol).
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

Read the thrombocytopenia tip sheet for more information.

Nausea and/or Vomiting

Take anti-nausea medications as prescribed. If you continue to have nausea or vomiting, notify your doctor or nurse so they can help you manage this side effect. In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try antacids, (e.g. milk of magnesia, calcium tablets such as Tums), saltines, or ginger ale to lessen symptoms. Read the Nausea & Vomiting Tip Sheet for more suggestions.

Call your doctor or nurse if you are unable to keep fluids down for more than 12 hours

or if you feel lightheaded or dizzy at any time.

Decrease in Appetite

Visit OncoLink's section on Nutrition for tips on dealing with this side effect. Ask your nurse about nutritional counseling services.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.
- You may experience a metallic taste or dislike foods or beverages that you liked before receiving chemotherapy. These symptoms can last up to several months.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products and fish without a strong smell.
- Flavor meat or fish by marinating it in sweet juices, sweet and sour sauce or dressings. Use seasonings like basil, oregano or rosemary. Bacon, ham and onion can also add flavor to vegetables.

Loss or Thinning of Scalp and Body Hair (Alopecia)

Your hair may become thin, brittle, or may fall out. This typically begins two to three weeks after treatment starts. This hair loss can be all body hair, including pubic, underarm, legs/arms, eyelashes, and nose hairs. The use of scarves, wigs, hats and hairpieces may help. Hair generally starts to regrow soon after treatment is completed. Remember your hair helps keep you warm in cold weather, so a hat is particularly important in cold weather or to protect you from the sun. Read more on alopecia.

Sexual and Reproductive Changes

This drug can affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and vaginal dryness - read more about coping with vaginal dryness. In addition, the desire for sex may decrease during treatment.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment, even if your menstrual cycle stops or you believe your sperm is affected. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team. See OncoLink's section on sexuality for helpful tips for dealing with these side effects.

Effect on Kidneys and Bladder

Ifosfamide can irritate the kidneys and damage the lining of the bladder, causing you to have blood in your urine. Intravenous fluids may be given with your treatment to flush the kidneys and bladder. A medication called Mesna may be given to protect the bladder. Drink 6-8 glasses of fluid a day. Urinate every 2-3 hours. Call your doctor or nurse if you have difficulty urinating or have blood in your urine.

Nervous System Toxicity

In rare cases, ifosfamide can be toxic to the nervous system. This can range from mild confusion, drowsiness, hallucinations and personality changes to seizure and coma. Toxicity can develop immediately after treatment up to several days later. Your caregiver should be aware of this rare side effect and know to report any changes in your behavior to your healthcare team right away.

Secondary Malignancies

There is a small risk of developing leukemia or other type of blood cancer due to treatment with this medication.

Cisplatin (Platinol®)

Pronounced: SIS-plat-in
Classification: Platinum Coordination Complex

About Cisplatin

Cisplatin is a heavy metal compound that inhibits synthesis of RNA, DNA, and protein in cells. All of these compounds are vital for cells to divide and grow. By preventing them from dividing, the medication can stop the cancer from growing.

How to Take Cisplatin

Cisplatin is given through intravenous (into a vein) infusion. The dose and schedule is determined by the person's size and type of cancer. It can be given alone or with other drugs.

Possible Side Effects of Cisplatin

There are a number of things you can do to manage the side effects of Cisplatin. Talk to your doctor or nurse about these recommendations. They can help you decide what will work best for you. These are some of the most common side effects:

Low White Blood Cell Count (Leukopenia or Neutropenia)

White blood cells (WBC) are important for fighting infection. While receiving treatment, your WBC count can drop, putting you at a higher risk of getting an infection. You should let your doctor or nurse know right away if you have a fever (temperature greater than 100.4 F), sore throat or cold, shortness of breath, cough, burning with urination, or a sore that doesn't heal.

Tips to preventing infection:

- Washing hands, both yours and your visitors, is the best way to prevent the spread of infection.
- Avoid large crowds and people who are sick (i.e.: those who have a cold, fever

or cough or live with someone with these symptoms).

- When working in your yard, wear protective clothing including long pants and gloves.
- Do not handle pet waste.
- Keep all cuts or scratches clean.
- Shower or bath daily and perform frequent mouth care.
- Do not cut cuticles or ingrown nails. You may wear nail polish, but not fake nails.
- Ask your doctor or nurse before scheduling dental appointments or procedures.
- Ask your doctor or nurse before you, or someone you live with, has any vaccinations.

For more suggestions, read the Neutropenia Tip Sheet.

Low Red Blood Cell Count (Anemia)

Your red blood cells are responsible for carrying oxygen to the tissues in your body. When the red cell count is low, you may feel tired or weak. You should let your doctor or nurse know if you experience any shortness of breath, difficulty breathing or pain in your chest. If the count gets too low, you may receive a blood transfusion. Read the anemia tip sheet for more information.

Low Platelet Count (Thrombocytopenia)

Platelets help your blood clot, so when the count is low you are at a higher risk of bleeding. Let your doctor or nurse know if you have any excess bruising or bleeding, including nosebleeds, bleeding gums or blood in your urine or stool. If your platelet count becomes too low, you may receive a transfusion of platelets.

- Do not use a razor to shave (an electric razor is fine).
- Avoid contact sports and activities that can result in injury or bleeding.
- Do not take aspirin (salicylic acid), non-steroidal, anti-inflammatory medications (NSAIDs) such as Motrin®, Aleve®, Advil®, etc. as these can all increase the risk of bleeding. Unless your healthcare team tells you otherwise, you may take acetaminophen (Tylenol).
- Do not floss or use toothpicks and use a soft-bristle toothbrush to brush your teeth.

Read the thrombocytopenia tip sheet for more information.

Nausea and/or Vomiting

Take anti-nausea medications as prescribed. If you continue to have nausea or vomiting, notify your doctor or nurse so they can help you manage this side effect. In addition, dietary changes may help. Avoid things that may worsen the symptoms, such as heavy or greasy/fatty, spicy or acidic foods (lemons, tomatoes, oranges). Try antacids, (e.g. milk of magnesia, calcium tablets such as Tums), saltines, or ginger ale to lessen symptoms. Read the Nausea & Vomiting Tip Sheet for more suggestions.

Call your doctor or nurse if you are unable to keep fluids down for more than 12 hours or if you feel lightheaded or dizzy at any time.

Decrease in Appetite

Visit OncoLink's section on Nutrition for tips on dealing with this side effect. Ask your nurse about nutritional counseling services.

- Try to eat five or six small meals or snacks throughout the day, instead of 3 larger meals.
- If you are not eating enough, nutritional supplements may help.
- You may experience a metallic taste or dislike foods or beverages that you liked before receiving chemotherapy. These symptoms can last up to several months.
- Avoid any food that you think smells or tastes bad. If red meat is a problem, eat chicken, turkey, eggs, dairy products and fish without a strong smell.
- Flavor meat or fish by marinating it in sweet juices, sweet and sour sauce or dressings. Use seasonings like basil, oregano or rosemary. Bacon, ham and onion can also add flavor to vegetables.

Sexual and Reproductive Changes

This drug can affect your reproductive system, resulting in the menstrual cycle or sperm production becoming irregular or stopping permanently. Women may experience menopausal effects including hot flashes and vaginal dryness – read more about coping with vaginal dryness. In addition, the desire for sex may decrease during treatment.

Exposure of an unborn child to this medication could cause birth defects, so you should not become pregnant or father a child while on this medication. Effective birth control is necessary during treatment, even if your menstrual cycle stops or you believe your sperm is affected. You may want to consider sperm banking or egg harvesting if you may wish to have a child in the future. Discuss these options with your oncology team. See OncoLink's section on sexuality for helpful tips for dealing with these side effects.

Kidney and Liver Function Changes

Your doctor may order some blood tests to check your liver and kidney function, as these can be affected by cisplatin. Call your doctor or nurse immediately if you have abdominal pain, if your urine becomes dark, or if your skin or the whites of your eyes become jaundiced (yellow in color).

Peripheral Neuropathy (Numbness or Tingling in the Hands and/or Feet)

Peripheral neuropathy is a toxicity that affects the nerves. It causes a numbness or tingling feeling in the hands and feet, often in the pattern of a stocking or glove. This can get progressively worse with additional doses of the drug. In some people, the symptoms slowly resolve after the drug is stopped, but for some it never goes away completely. You should let your healthcare provider know if you experience numbness or tingling in the hands and feet, as they may need to change the doses of your medication. See OncoLink's section on peripheral neuropathy for tips on dealing with this side effect.

Electrolyte Changes

Your doctor may order blood tests to check your electrolytes. If needed, the doctor may prescribe supplements such as magnesium or potassium.

Hearing Loss

Cisplatin can cause hearing loss. Call your doctor or nurse if you have ringing in your ears or if you notice a decrease in your hearing.

© Trustees of the University of Pennsylvania